

# AllBio Xba I-Fast Cut



Cat. No. ABTGREX101

Storage : at -20°C for two years

Concentration : 20,000 units/ml

Recognition Site

5'...T<sup>▼</sup>CTAGA...3'

3'...AGATCT<sup>▲</sup>...5'

## Description

AllBio Xba I-Fast Cut is expressed and purified from E.coli that carries the recombinant Xba I gene. The molecular weight is 24.7 kDa, with the recognition site at T<sup>▼</sup>CTAGA. The reaction is conducted for 5-15 minutes at 37°C, and heat-inactivated at 65°C for 20 minutes. This enzyme is not sensitive to dcm or mammalian CpG methylation, but sensitive to dam methylation.

## Enzyme Properties

Fast digestion in 5-15 minutes with high fidelity

## Application

Genomic DNA, plasmid DNA, PCR product

## Kit Contents

Component	ABTGREX101-01	ABTGREX101-02
AllBio Xba I-Fast Cut	1,500 units	2×1,500 units
10×FastCut Buffer	500 µl	1 ml
10×DNA Loading Buffer	1 ml	1 ml

## Unit Definition

One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

## Quality Control

**Ligation and re-cutting:** After 10-fold overdigestion with AllBio Xba I-Fast, more than 95% of the DNA fragments can be ligated with T4 DNA ligase at 25°C. Of these ligated fragments, more than 95% can be recut.

**16-Hour incubation:** A 50 µl reaction containing 1 µg of DNA and 10 units of enzyme incubated for 16 hours results in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Blue/White screening (Terminal integrity):** A DNA vector is digested at a unique site within lacZα gene with a 10-fold excess of enzyme, and then ligated, transformed and plated on X-gal/IPTG plate. Successful expression of the β-galactosidase indicates that lacZα gene remains integrity after cloning. A blue colony represents an intact gene, and a white colony represents an interrupted gene. To be Blue/White certified, enzymes must produce fewer than 3% white colonies.

**Exonuclease activity:** After incubation for 4 hours at 37°C, a 50 µl reaction containing 100 units of enzyme and 1 µg <sup>3</sup>H DNA releases less than 0.1% radioactive substance.

**Endonuclease activity:** After incubation for 4 hours at 37°C, a 50 µl reaction containing 15 units of enzyme with 1 µg pBR322 RFI DNA results in less than 10% conversion from RFI to RFII.

## Storage Buffer

20 mM Tris-HCl pH7.4, 250 mM NaCl, 0.1 mM EDTA, 1.5 mM DTT, 400 µg/ml BSA, 50% Glycerol

## 10×FastCut Buffer

500 mM Tris-Ac pH7.9, 1 M KAc, 120 mM MgAc<sub>2</sub>, 1 mg/ml BSA

# AllBio Xba I-Fast Cut



## Reaction Components

Component	20 $\mu$ l Volume	50 $\mu$ l Volume
DNA	$\leq 1 \mu\text{g}$	$\leq 2 \mu\text{g}$
10 $\times$ FastCut Buffer	2 $\mu$ l	5 $\mu$ l
AllBio Xba I-Fast Cut	0.5 $\mu$ l	1 $\mu$ l
ddH <sub>2</sub> O	to 20 $\mu$ l	to 50 $\mu$ l

For digestion of  $>2 \mu\text{g}$  DNA, please adjust the amount of each component based on above table.

## Reaction Condition

Incubation for 5-10 minutes at 37°C. Enzyme is inactivated by adding 10 $\times$ DNA Loading Buffer to a final concentration at 1 $\times$ , or by heating at 65°C for 20 minutes.

## Notes

- Thaw the 10 $\times$ FastCut Buffer completely and mix well before use.
- Low ionic strength, high enzyme concentration, glycerol concentration  $> 5\%$ , or pH  $> 8.0$  may result in star activity.